

# PATENT COOPERATION TREATY

## PCT

### INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference <b>80021-210</b>	<b>FOR FURTHER ACTION</b> see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.	
International application No. <b>PCT/CA 00/ 00860</b>	International filing date (day/month/year) <b>21/07/2000</b>	(Earliest) Priority Date (day/month/year) <b>22/07/1999</b>
Applicant  <b>THE UNIVERSITY OF BRITISH COLUMBIA</b>		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.



It is also accompanied by a copy of each prior art document cited in this report.

**1. Basis of the report**

a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.



the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :



contained in the international application in written form.



filed together with the international application in computer readable form.



furnished subsequently to this Authority in written form.



furnished subsequently to this Authority in computer readable form.



the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.



the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☐ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (see Box II).

4. With regard to the **title**,



the text is approved as submitted by the applicant.



the text has been established by this Authority to read as follows:

5. With regard to the **abstract**,



the text is approved as submitted by the applicant.



the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

6. The figure of the **drawings** to be published with the abstract is Figure No.



as suggested by the applicant.



because the applicant failed to suggest a figure.



because this figure better characterizes the invention.



None of the figures.

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/CA 00/00860

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/54 C12N15/82 C12N9/10 A01H5/00 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A01H C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, STRAND, MEDLINE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EMBL 'Online! Acc no. ATT4L20, 29 April 1998 (1998-04-29) BEVAN M. ET AL: "Arabidopsis thaliana DNA chromosome 4, BAC clone T4L20 (ESSA project)" XP002162955 nts 26761-29269	1-37
X	WO 95 15387 A (LARDIZABAL KATHRYN DENNIS ; LASSNER MICHAEL W (US); METZ JAMES GEOR) 8 June 1995 (1995-06-08) cited in the application SEQ ID NO.21	20-23, 34-37

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*G\* document member of the same patent family

Date of the actual completion of the international search

16 March 2001

Date of mailing of the international search report

28/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+31-70) 340-3016

Authorized officer

Blanco Urgoiti, B

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 00/00860

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	TODD JAMES ET AL: "KCS1 encodes a fatty acid elongase 3-ketoacyl-CoA synthase affecting wax biosynthesis in Arabidopsis thaliana." PLANT JOURNAL, vol. 17, no. 2, January 1999 (1999-01), pages 119-130, XP002162954 ISSN: 0960-7412	
A	WO 96 13582 A (DNA PLANT TECHN CORP) 9 May 1996 (1996-05-09) cited in the application the whole document	

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
1 February 2001 (01.02.2001)

PCT

(10) International Publication Number  
WO 01/07586 A2

- (51) International Patent Classification<sup>7</sup>: C12N 15/00 (74) Agents: KINGWELL, Brian, G. et al.; Smart & Biggar, Box 11560, Vancouver Centre, 650 West Georgia Street, Suite 2200, Vancouver, British Columbia V6B 4N8 (CA).
- (21) International Application Number: PCT/CA00/00860
- (22) International Filing Date: 21 July 2000 (21.07.2000) (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/145,013 22 July 1999 (22.07.1999) US (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): THE UNIVERSITY OF BRITISH COLUMBIA [CA/CA]; University-Industry Liaison Office, IRC Building, Room 331, 2194 Health Sciences Mall, Vancouver, British Columbia V6T 1Z3 (CA).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): KUNST, Ljerka [CA/CA]; 2810 Newmarket Drive, North Vancouver, British Columbia V7R 2T4 (CA). CLEMENS, Sabine [DE/CA]; 4443 West 16th Avenue, Vancouver, British Columbia V6R 3E7 (CA).

Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: A PLANT LONG CHAIN FATTY ACID BIOSYNTHETIC ENZYME

1 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
21 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
31 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
41 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
51 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
61 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
71 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
81 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
91 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
101 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
111 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
121 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
131 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
141 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
151 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
161 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
171 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
181 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
191 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
201 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
211 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
221 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
231 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
241 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
251 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
261 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
271 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
281 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
291 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
301 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
311 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
321 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
331 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
341 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
351 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
361 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
371 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
381 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
391 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
401 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
411 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
421 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
431 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
441 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
451 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
461 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
471 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
481 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
491 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
501 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
511 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
521 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
531 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
541 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
551 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
561 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
571 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
581 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
591 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
601 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
611 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
621 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
631 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
641 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
651 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
661 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
671 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
681 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
691 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
701 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
711 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
721 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
731 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
741 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
751 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
761 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
771 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
781 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
791 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
801 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
811 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
821 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
831 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
841 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
851 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
861 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
871 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
881 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
891 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
901 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
911 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
921 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
931 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
941 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
951 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
961 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
971 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
981 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
991 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG  
1001 TTTTGGGG GGGGGGGG GGGGGGGG GGGGGGGG GGGGGGGG

(57) Abstract: In various aspects, the present invention provides nucleic acid sequence encoding all or part of a new plant long chain fatty acid condensing enzyme (a fatty acid elongase), designated herein as KCS2 (for beta-ketoacyl-coenzyme A synthase 2). In some embodiments, KCS2 may mediate the biosynthesis of C18, C20, C22 and C24 fatty acids. The activity of the enzyme is typically characterized by two carbon (malonyl-CoA) additions to C16, C18, C20 and C22 moieties (C16-C22 acyl CoA molecules), i.e. condensation of malonyl-CoA with a C16, C18, C20 or C22 acyl-CoA. The fatty acids produced by the enzyme may for example be saturated 18:0, 20:0, 22:0 and 24:0 fatty acids. The invention includes recombinant nucleic acid molecules comprising a heterologous nucleic acid coding sequence encoding the plant long chain fatty acid condensing enzyme.

WO 01/07586 A2

## A PLANT LONG CHAIN FATTY ACID BIOSYNTHETIC ENZYME

### FIELD OF THE INVENTION

The invention relates to a condensing enzyme involved in long chain fatty acid  
5 production in plants, including related nucleic acid sequences.

### BACKGROUND

Living organisms synthesize a vast array of different fatty acids, which are incorporated  
into complex lipids. These complex lipids represent both major structural membrane  
10 components, and are a major storage product in both plants and animals. In plants, very long  
chain fatty acids (VLCFAs, chain length C20 or longer) are synthesized predominantly in the  
epidermal cells where they are either directly incorporated into waxes, or serve as precursors  
for other aliphatic hydrocarbons found in waxes, including alkanes, primary and secondary  
alcohols, ketones, aldehydes and acyl-esters (for review see Post-Beittenmiller, 1996).

15 VLCFAs also accumulate in the seed oil of some plant species, where they are incorporated  
into triacylglycerols (TAGs), as in the *Brassicaceae*, or into wax esters, as in Jojoba. These  
seed VLCFAs include the agronomically important erucic acid (C22:1), that may be used in  
the production of lubricants, nylon, cosmetics, pharmaceuticals and plasticizers.

20 VLCFAs are synthesized by a microsomal fatty acid elongation (FAE) system which  
involves four enzymatic reactions: (1) condensation of malonyl-CoA with a long chain acyl-  
CoA, (2) reduction to beta-hydroxyacyl-CoA, (3) dehydration to an enoyl-CoA and (4)  
reduction of the enoyl-CoA, resulting in the acyl-CoA elongated by two carbons. The  
condensing enzyme catalyzing reaction (1) is a key activity of the FAE system. It is the rate-  
25 limiting enzyme of the VLCFA biosynthetic pathway, which controls the amount of VLCFAs  
produced (Miller and Kunst, 1997). In addition, the condensing enzyme determines the  
ultimate VLCFA acyl chain length, and thus their uses in seed oil or wax biosynthesis.

30 Different condensing enzymes acting on, and producing, acyl chains of different  
length have recently been characterized. Several groups independently identified the first  
plant fatty acid elongation gene in *Arabidopsis*, *FAE1* (James and Dooner, 1990; Kunst et  
al., 1992; and Lemieux et al., 1990). *FAE1* was subsequently cloned and is described in  
WO9613582 as catalyzing the conversion of C18 fatty acids to C20-C22 fatty acids. The  
patent WO9613582 suggests that *FAE1* will have activity in a very broad host range. In

support of this assertion of broad host range activity, it has been shown that *FAE1* has the same activity in yeast as in *Arabidopsis* (Miller and Kunst, 1997). A Jojoba protein involved in the synthesis of VLCFAs has also been isolated having relatively high homology to *FAE1* (52% amino acid identity), and has been shown to have beta-ketoacyl-coenzyme A synthase (*KCS*) activity (WO9515387). Broad host range activity for genes encoding *KCS* has been further evidenced by the Jojoba *KCS* cDNA. The Jojoba *KCS* cDNA was able to complement a mutation in a Canola variety of rapeseed (*Brassica napus*), to restore in the variety high levels of VLCFAs (Lassner *et al.*, 1996). An *Arabidopsis* gene (*CUT1*), required for cuticular wax biosynthesis and pollen fertility, has also been described as encoding a VLCFA condensing enzyme that catalyzes the addition of 2C units to pre-existing C24 or longer fatty acids (Millar *et al.*, 1999).

The broad specificities exhibited by *FAE* activities provide a means of modifying the synthesis of VLCFAs in a given cell. As evidenced by the accumulation of VLCFAs in tobacco seed expressing *FAE1* (Miller and Kunst, 1997), heterologous condensing enzymes may be used to produce VLCFAs in plant species that do not otherwise synthesize VLCFAs. For example, targeted expression of heterologous VLCFA condensing enzymes in seeds may allow the production of crop plants capable of synthesizing VLCFAs of desired lengths in seed oil for industrial applications. New condensing enzymes may also be useful for the manipulation of cuticular waxes which have important functions in many physiological processes in plants, including water balance, protection from UV light, plant-insect interactions, and defense against bacterial and fungal pathogens (Post-Beittenmiller, 1996). As a result, there is a need for new condensing enzymes that may be used alone or in combination with other condensing enzymes to confer on plants and plant tissues the ability to synthesize a range of VLCFAs, including VLCFAs up to C30 in length.

There is also a need for tissue-specific promoters capable of mediating the expression of heterologous condensing enzymes in epidermal cells, which may facilitate the alteration of the wax composition and/or accumulation in plants. This may, in turn, result in the production of crops with increased tolerance to environmental stresses, and/or resistance to pathogens and insects. For example, drought resistance in rice is associated with high wax lines rich in C29, C33 and C35 alkanes (O'Toole and Cruz, 1983; Haque *et al.*, 1992). Increased wax deposition may also be accomplished by overexpression of condensing enzymes with desired acyl chain

length specificities using an epidermis-specific promoter, such as the *CUT1* promoter (Millar et al., 1999).

The cumulative data, as discussed, from a variety of sources in this field has led to the suggestion that the amounts and acyl chain lengths of VLCFAs, in a wide variety of eukaryotic cells, are regulated by the nature of condensing enzyme expression in the cell (Miller and Kunst, 1997). Condensing enzymes would therefore be useful in a range of biotechnical applications.

## 10 SUMMARY

In various aspects, the present invention provides nucleic acid sequence encoding all or part of a new plant long chain fatty acid condensing enzyme (fatty acid elongase), designated herein as KCS2 (for beta-ketoacyl-coenzyme A synthase 2). In some embodiments, KCS2 may mediate the biosynthesis of C18, C20, C22 and C24 fatty acids. The activity of the enzyme is typically characterized by two carbon (malonyl-CoA) additions to C16, C18, C20 and C22 moieties (C16-C22 acyl CoA molecules), i.e. condensation of malonyl-CoA with a C16, C18, C20 or C22 acyl-CoA. The fatty acids produced by the enzyme may for example be saturated 18:0, 20:0, 22:0 and 24:0 fatty acids.

The invention includes recombinant nucleic acid molecules comprising a heterologous nucleic acid coding sequence encoding the plant long chain fatty acid condensing enzyme. In alternative embodiments, the nucleic acid coding sequence may be derived from the *Arabidopsis* KCS2 coding sequence disclosed herein. Alternatively, embodiments include nucleic acids that encode the plant very long chain fatty acid condensing enzyme of the invention, wherein the enzyme has an amino acid sequence that is at least 70% identical to an *Arabidopsis* KCS2 amino acid sequence disclosed herein, when optimally aligned. The nucleic acid coding sequences of the invention also include sequences that hybridize under stringent conditions to a complement of the *Arabidopsis* KCS2 coding sequence disclosed herein. The nucleic acid coding sequences of the invention may also be at least 70% identical to the *Arabidopsis* KCS2 coding sequence, when optimally aligned. Embodiments of the invention include isolated nucleic acid molecules comprising the coding sequences of the invention.

In another aspect, the invention provides recombinant nucleic acid molecules comprising a promoter sequence operably linked to a nucleic acid sequence, wherein the

promoter sequence is capable of mediating gene expression in anthers and in very young leaves in *Arabidopsis*. The promoter sequences of the invention may be derived from an *Arabidopsis KCS2* promoter sequence, as disclosed herein. Promoter sequences of the invention may also hybridize under stringent conditions to the *Arabidopsis KCS2* promoter sequence disclosed  
5 herein. Promoter sequences of the invention may also be at least 70% identical to the *Arabidopsis KCS2* promoter sequence when optimally aligned.

The invention provides nucleic acid probes comprising probe sequences that hybridize under stringent conditions to a portion of an *Arabidopsis KCS2* genomic sequence; or are at  
10 least 70% identical to the portion of an *Arabidopsis KCS2* genomic sequence when optimally aligned. The invention also provides methods of isolating a nucleic acid molecule encoding a plant long chain fatty acid condensing enzyme, for example by hybridizing a nucleic acid preparation with the nucleic acid probe of the invention.

The invention provides transgenic cells (such as plant cells), plants and parts thereof, in  
15 which the plants or cells comprise the recombinant nucleic acid molecules of the invention. Such plant parts may for example include seeds or oils. Transgenic plants of the invention may have a modified phenotype compared to a non-transgenic plant of the same species, such as a modified lipid content. Methods of producing such transgenic plants are provided, for example  
20 by introducing into a plant the isolated nucleic acids of the invention. Progeny plants may be provided, produced by sexual or asexual propagation of the transgenic plants of the invention to produce transgenic descendants of transformed plants.

Purified proteins are provided, encoded by the recombinant nucleic acid molecules of  
25 the invention, including plant fatty acid condensing (elongase) enzymes and fragments thereof. Also provided are recombinant vectors comprising the recombinant nucleic acid molecules of the invention.

Antisense nucleic acid molecules are provided, wherein a portion of the nucleic acid  
30 coding sequences of the invention are provided in reverse orientation relative to an adjacent promoter sequence. Recombinant antisense nucleic acids of the invention may therefore encode an antisense RNA that hybridizes under stringent conditions to a complement of a portion of the *Arabidopsis KCS2* coding sequence; or that are at least 70% identical to a portion of the *Arabidopsis KCS2* coding sequence when optimally aligned. Transgenic plants



or plant cells of the invention may include the recombinant antisense nucleic acids of the invention.

In one embodiment, the nucleic acid coding sequences of the invention may be substantially identical to all or part of an *Arabidopsis KCS2* coding sequence. The nucleic acids of the invention may also include an RNA analog or a nucleic acid complementary to sequences of the invention. In other embodiments, the nucleic acid may be a fragment of one of the above sequences, such as a fragment that is at least 5, 10, 15, 20 or 25 nucleotides in length and that hybridizes under stringent conditions to a genomic *KCS2* DNA encoding the nucleic acid sequence. As used herein, the term "genomic sequence" includes either of the strands of a nucleic acid molecule found in the genome of an organism or cell.

In another aspect, the nucleic acids of the invention may include a DNA coding sequence encoding an enzyme of the invention, operably linked to a promoter. Promoters of the invention may be tissue-specific or have specific developmental timing. A promoter of the invention may have a nucleotide sequence substantially identical to all or part of the *KCS2* promoter region disclosed herein. Promoters of the invention may be operably linked to alternative DNAs, such as agronomically important nucleic acid sequences.

In one aspect, the invention provides methods for altering the VLCFA, fatty acid or lipid content in a plant or plant tissue, for example by introducing into a plant cell, capable of being transformed and regenerated to a whole plant, a nucleic acid of the invention. Where such nucleic acids are effective for altering the levels of VLCFAs in a plant; a plant containing the nucleic acid of the invention may be recovered having an altered phenotype.

## THE DRAWINGS

Figure 1 shows the coding strand DNA sequence of a *KCS2* genomic clone from *Arabidopsis*, containing a 1001 bp long *KCS2* promoter region (5' to the initiation codon) and a 1508 bp long *KCS2* coding sequence. The initiation (ATG) codon is indicated in all caps (at positions 1046-1048). The Figure identifies locations homologous or complementary to oligonucleotide primers (probes) that may be used for amplifying the wild-type *Arabidopsis KCS2* promoter (shown as *kcs3* and *kcs4*) and the *KCS2* coding sequence (*kcs1* and *kcs2*), these sequences are underlined and marked by arrows indicating the putative direction for amplification.

## DETAILED DESCRIPTION OF THE INVENTION

The recombinant nucleic acid molecules of the invention may comprise a heterologous nucleic acid coding sequence encoding the plant long chain fatty acid condensing enzyme of the invention. In alternative embodiments, the nucleic acid coding sequence may be derived from the *Arabidopsis KCS2* coding sequence disclosed herein.

The term "recombinant" means that something has been recombined, so that when made in reference to a nucleic acid molecule the term refers to a molecule that is comprised of nucleic acid sequences that are joined together by means of molecular biological techniques. The term "recombinant" when made in reference to a protein or a polypeptide refers to a protein molecule which is expressed using a recombinant nucleic acid molecule. The term "heterologous" when made in reference to a nucleic acid sequence refers to a nucleotide sequence which is ligated to, or is manipulated to become ligated to, a nucleic acid sequence to which it is not ligated in nature, or to which it is ligated at a different location in nature. The term "heterologous" therefore indicates that the nucleic acid molecule has been manipulated using genetic engineering, i.e. by human intervention.

Sequences may be derived or obtainable from the *Arabidopsis KCS2* coding sequence by deduction and synthesis based upon the wild-type KCS2 amino acid sequence, using the redundancy of the genetic code to formulate alternative coding sequences. Derived sequences may also be identified in different organisms, for example by isolation using as probes the nucleic acid sequences of the invention. Antibodies prepared against a KCS2 protein of the invention may also be used to identify alternative coding sequences, which are thereby derived from the coding sequence disclosed herein. Alternative KCS2 amino acid sequences of the invention, such as amino acid sequences developed through mutagenesis or substitution, may similarly be used to infer coding sequences derived from the *Arabidopsis KCS2* coding sequence disclosed herein.

Derived nucleic acids of the invention may be obtained by chemical synthesis, isolation, or cloning from genomic DNAs using techniques known in the art, such as the Polymerase Chain Reaction (PCR). Nucleic acids of the present invention may be used to design alternative primers (probes) suitable for use as PCR primers to amplify particular regions of an condensing enzyme cDNA of the invention. Such PCR primers may for example

comprise a sequence of 15-20 consecutive nucleotides of the KCS2 cDNA of the invention. To enhance amplification specificity, primers of 20-30 nucleotides in length may also be used. Methods and conditions for PCR amplification are described in Innis et al. (1990); Sambrook et al. (1989); and Ausubel et al. (1995).

5

As used herein, the term "probe" when made in reference to an oligonucleotide refers to an oligonucleotide which is capable of hybridizing to another oligonucleotide of interest. A probe may be single-stranded or double-stranded. Probes are, for example, useful in the detection, identification, amplification and isolation of particular gene sequences.

10 Oligonucleotide probes may be labelled with a "reporter molecule," so that the probe is detectable using a detection system, such as enzymatic, fluorescent, radioactive or luminescent detection systems.

Derived nucleic acids of the invention may also be identified by Southern analysis, a  
15 method by which the presence of DNA sequences in a target nucleic acid mixture are identified by hybridization to a labeled probe, comprising an oligonucleotide or DNA fragment of a nucleic acid of the invention. Probes for Southern analysis may for example be at least 15 nucleotides in length. Southern analysis typically involves electrophoretic separation of DNA digests on agarose gels, denaturation of the DNA after electrophoretic separation, and transfer  
20 of the DNA to nitrocellulose, nylon, or another suitable membrane support for analysis with a radiolabeled, biotinylated, or enzyme-labeled probe as described in Sambrook *et al.* (1989).

In alternative embodiments, the invention includes nucleic acids that encode the plant very long chain fatty acid condensing enzyme of the invention, wherein the enzyme has an  
25 amino acid sequence that is at least 70% identical to an *Arabidopsis* KCS2 amino acid sequence disclosed herein, when optimally aligned. In alternative embodiments, the degree of identity may be between 50% and 100%, such as 60%, 80%, 90%, 95% or 99%.

As is well known in the art, some modifications and changes can be made in the  
30 structure of a polypeptide without substantially altering the biological function of that peptide, to obtain a biologically equivalent polypeptide. In one aspect of the invention, fatty acid condensing enzymes may include peptides that differ from a portion of the wild-type *Arabidopsis* KCS2 sequence by conservative amino acid substitutions. As used herein, the term "conserved amino acid substitutions" refers to the substitution of one amino acid for another at

a given location in the peptide, where the substitution can be made without substantial loss of function. In making such changes, substitutions of like amino acid residues can be made, for example, on the basis of relative similarity of side-chain substituents, for example, their size, charge, hydrophobicity, hydrophilicity, and the like, and such substitutions may be assayed for their effect on the function of the peptide by routine testing.

In some embodiments, conserved amino acid substitutions may be made where an amino acid residue is substituted for another having a similar hydrophilicity value (e.g., within a value of plus or minus 2.0), where the following hydrophilicity values are assigned to amino acid residues (as detailed in United States Patent No. 4,554,101, incorporated herein by reference): Arg (+3.0); Lys (+3.0); Asp (+3.0); Glu (+3.0); Ser (+0.3); Asn (+0.2); Gln (+0.2); Gly (0); Pro (-0.5); Thr (-0.4); Ala (-0.5); His (-0.5); Cys (-1.0); Met (-1.3); Val (-1.5); Leu (-1.8); Ile (-1.8); Tyr (-2.3); Phe (-2.5); and Trp (-3.4).

In alternative embodiments, conserved amino acid substitutions may be made where an amino acid residue is substituted for another having a similar hydropathic index (e.g., within a value of plus or minus 2.0). In such embodiments, each amino acid residue may be assigned a hydropathic index on the basis of its hydrophobicity and charge characteristics, as follows: Ile (+4.5); Val (+4.2); Leu (+3.8); Phe (+2.8); Cys (+2.5); Met (+1.9); Ala (+1.8); Gly (-0.4); Thr (-0.7); Ser (-0.8); Trp (-0.9); Tyr (-1.3); Pro (-1.6); His (-3.2); Glu (-3.5); Gln (-3.5); Asp (-3.5); Asn (-3.5); Lys (-3.9); and Arg (-4.5).

In alternative embodiments, conserved amino acid substitutions may be made where an amino acid residue is substituted for another in the same class, where the amino acids are divided into non-polar, acidic, basic and neutral classes, as follows: non-polar: Ala, Val, Leu, Ile, Phe, Trp, Pro, Met; acidic: Asp, Glu; basic: Lys, Arg, His; neutral: Gly, Ser, Thr, Cys, Asn, Gln, Tyr.

The nucleic acid coding sequences of the invention also include sequences that hybridize under stringent conditions to a complement of the *Arabidopsis KCS2* coding sequence disclosed herein. The nucleic acid coding sequences of the invention may also be at least 70% identical to the *Arabidopsis KCS2* coding sequence, when optimally aligned. Embodiments of the invention include isolated nucleic acid molecules comprising the coding sequences of the invention.

The present invention provides an isolated nucleic acid encoding a plant elongase enzyme that mediates biosynthesis of 18:0, 20:0, 22:0 and 24:0 fatty acids in plants. In a preferred embodiment, the nucleic acid encodes *KCS2* identified in *Arabidopsis* shown in Figure 1.

By isolated, it is meant that the isolated substance has been substantially separated or purified away from other biological components with which it would otherwise be associated, for example *in vivo*. The term 'isolated' therefore includes substances purified by standard purification methods, as well as substances prepared by recombinant expression in a host, as well as chemically synthesized substances.

The nucleic acids of this invention may be genomic or cDNA and may be isolated from cDNA or genomic libraries or directly from isolated plant DNA. Nucleic acids of this invention further include sequences corresponding to the *KCS2* protein as well as sequences obtainable from the *KCS2* protein or nucleic acid sequences. By corresponding is meant nucleic acid sequences, either DNA or RNA, including those which encode the *KCS2* protein or a portion thereof, the promoter sequence found 5' to said encoding sequence, intron sequences not present in the cDNA, as well as sequences encoding any leader or signal peptide of a precursor protein that may not be found in the mature elongase protein. A nucleic acid of the invention may further comprise additional nucleic acids. For example, linkers, modified or unmodified restriction endonuclease sites and other nucleic acid sequences useful for cloning, expression, or purification.

Sequences having substantial identity will hybridize under stringent conditions. Hybridization to filter-bound sequences may for example, be performed in 0.5 M NaHPO<sub>4</sub>, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.2 x SSC/0.1% SDS at 42°C (Ausubel *et al.*, 1995). Alternatively hybridization to filter-bound sequences may for example, be performed in 0.5 M NaHPO<sub>4</sub>, 7% sodium dodecyl sulfate (SDS), 1 mM EDTA at 65°C, and washing in 0.1 x SSC/0.1% SDS at 68°C (Ausubel *et al.*, 1995). Hybridization conditions may be modified in accordance with known methods depending on the sequence of interest (Tijssen, 1993). Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point for the specific sequence at the defined ionic strength and pH.

A nucleic acid may also be identified by Northern analysis, a method used to identify RNAs that hybridize to a known probe such as an oligonucleotide, DNA fragment, cDNA or fragment thereof, or RNA fragment of a nucleic acid of the invention. The probe is labeled with a radioisotope such as  $^{32}\text{P}$ , by biotinylation or with an enzyme. The RNA to be analyzed  
5 may be electrophoretically separated on an agarose or polyacrylamide gel, transferred to nitrocellulose, nylon, or other suitable membrane, and hybridized with the probe, using standard techniques well known in the art such as described in Sambrook *et al.* (1989).

A nucleic acid of the invention may alternatively be obtained by immunological  
10 screening. Antibodies to the KCS2 protein may be used to identify related protein in extracts of desired plant species. Two amino acid sequences are considered substantially identical if one peptide is specifically immunologically reactive with antibodies that are also specifically immunoreactive against the other peptide. Specific immunoreactivity of antibodies to peptides may be assessed using a variety of immunoassay formats, such as solid-phase ELISA  
15 immunoassays for selecting monoclonal antibodies specifically immunoreactive with a protein (Harlow and Lane, 1988).

Optimal alignment of sequences for comparisons of identity may be conducted using a variety of algorithms, such as the local homology algorithm of Smith and Waterman (1981), the homology alignment algorithm of Needleman and Wunsch (1970), the search for similarity  
20 method of Pearson and Lipman (1988), and the computerized implementations of these algorithms (such as GAP, BESTFIT, FASTA and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, Madison, WI, U.S.A.). Sequence alignment may also be carried out using the BLAST algorithm described in Altschul *et al.* (1990), using the published default settings.

25 Software for performing BLAST analysis may be available through the National Center for Biotechnology Information (through the internet at <http://www.ncbi.nlm.nih.gov/>). The BLAST algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length  $W$  in the query sequence that either match or satisfy some positive-valued threshold score  $T$  when aligned with a word of the same length in a database sequence.  
30  $T$  is referred to as the neighbourhood word score threshold. Initial neighbourhood word hits act as seeds for initiating searches to find longer HSPs. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Extension of the word hits in each direction is halted when the following parameters are met:

the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLAST programs may use as defaults a word length (W) of 11, the BLOSUM62 scoring matrix (Henikoff and Henikoff, 1992, *Proc. Natl. Acad. Sci. USA* 89: 10915-10919) alignments (B) of 50, expectation (E) of 10 (which may be changed in alternative embodiments to 1 or 0.1 or 0.01 or 0.001 or 0.0001; although E values much higher than 0.1 may not identify functionally similar sequences, it is useful to examine hits with lower significance, E values between 0.1 and 10, for short regions of similarity), M=5, N=4, for nucleic acids a comparison of both strands. For protein comparisons, BLASTP may be used with defaults as follows: G=11 (cost to open a gap); E=1 (cost to extend a gap); E=10 (expectation value, at this setting, 10 hits with scores equal to or better than the defined alignment score, S, are expected to occur by chance in a database of the same size as the one being searched; the E value can be increased or decreased to alter the stringency of the search.); and W=3 (word size, default is 11 for BLASTN, 3 for other blast programs).

The BLOSUM matrix assigns a probability score for each position in an alignment that is based on the frequency with which that substitution is known to occur among consensus blocks within related proteins. The BLOSUM62 (gap existence cost = 11; per residue gap cost = 1; lambda ratio = 0.85) substitution matrix is used by default in BLAST 2.0. A variety of other matrices may be used as alternatives to BLOSUM62, including: PAM30 (settings: 9,1,0.87); PAM70 (settings: 10,1,0.87) BLOSUM80 (settings: 10,1,0.87); BLOSUM62 (settings: 11,1,0.82) and BLOSUM45 (settings: 14,2,0.87). One measure of the statistical similarity between two sequences using the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. In alternative embodiments of the invention, nucleotide or amino acid sequences may be identified as sequences of the invention where the smallest sum probability in a comparison to a reference KCS2 amino acid or KCS2 nucleic acid test sequence is less than about 1, preferably less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

When a position in the compared sequence is occupied by the same nucleotide or amino acid, following optimal alignment of the sequences, the molecules are considered to have

identity at that position. The degree of identity between sequences is a function of the number of matching positions shared by the sequences. In terms of percentage, identity is the sum of identical positions, divided by the total length over which the sequences are aligned, multiplied by 100.

5

It will be recognized by one of ordinary skill in the art that nucleic acids of this invention may be modified using standard techniques of site specific mutation or PCR, or modification of the sequence may be accomplished in producing a synthetic nucleic acid sequence. Such modified sequences are also considered in this invention. For example, due to the degeneracy of the genetic code, which is well-known to the art; i.e., for many amino acids, there is more than one nucleotide triplet which serve as the codon for the amino acid, codons may be changed such that the nucleic acid sequence encodes the same amino acid sequence, or alternatively, codons may be altered such that conservative amino acid substitutions or substitutions of similar amino acids result without affecting protein function.

15

In another aspect of the invention, the nucleic acid may further comprise a promoter operably linked to the enzyme-encoding region. A first nucleic acid sequence is operably linked with a second nucleic acid sequence when the first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequences. Generally, operably linked DNA sequences are contiguous and, where necessary to join two protein coding regions, in reading frame.

A promoter, as used herein, is a DNA sequence in a gene, usually (but not necessarily) upstream (5') to its coding sequence, which controls the expression of the coding sequence by providing the recognition for RNA polymerase and other factors required for proper transcription. In artificial DNA constructs promoters may also be used to transcribe antisense RNA. Promoters may also contain DNA sequences that are involved in the binding of protein factors which control the effectiveness of transcription initiation in response to physiological or developmental conditions. A promoter may also contain enhancer elements, a DNA sequence which can stimulate promoter activity. A gene of the present invention may also include transcription termination signals.

30



In one embodiment of the invention, the nucleic acid may be used to alter the composition and accumulation of fatty acids in plant cells. For example, for over-expression, which refers to production of a gene product exceeding levels of production in normal or non-transformed organisms, a plant promoter may be employed which will direct expression of the  
 5 elongase in all tissues of a regenerated plant. Over-expression of the gene, for example, in plant epidermal cells could increase cuticle accumulation thereby increasing drought and stress tolerance of transgenic plants over control plants. Such promoters are referred to herein as “constitutive” promoters and are active under most environmental conditions and states of development or cell differentiation. Examples of constitutive promoters include the  
 10 cauliflower mosaic virus (CaMV) 35S transcription initiation region, the 1’- or 2’- promoter derived from T-DNA of *Agrobacterium tumefaciens*, and other transcription initiation regions from various plant genes known to those of skill in the art.

In various embodiments, the invention comprises plants transformed with the nucleic  
 15 acids of the invention. In some embodiments, such plants will exhibit altered fatty acid content in one or more tissues. These aspects of the invention relate to all higher plants, including monocots and dicots, such as species from the genera *Fragaria*, *Lotus*, *Medicago*, *Onobrychis*, *Trifolium*, *Trigonelia*, *Wigna*, *Citrus*, *Linum*, *Geranium*, *Manihot*, *Caucus*, *Arabidopsis*, *Brassica*, *Raphanus*, *Sinapis*, *Atropa*, *Capsicum*, *Hyoscyamus*, *Lycopersicon*, *Nicotiana*,  
 20 *Solanum*, *Petunia*, *Digitalis*, *Majorana*, *Cichorium*, *Helianthus*, *Lactuca*, *Bromus*, *Asparagus*, *Antirrhinum*, *Heterocatlis*, *Nemesia*, *Pelargonium*, *Panicum*, *Pennisetum*, *Ranunculus*, *Senecio*, *Salpiglossis*, *Cucumis*, *Browallia*, *Glycine*, *Lolium*, *Zea*, *Triticum*, *Sorghum*, and *Datura*. Such plants may include maize, wheat, rice, barley, soybean, beans, rapeseed, canola, alfalfa, flax, sunflower, cotton, clover, lettuce, tomato cucurbits, potato carrot, radish, pea  
 25 lentils, cabbage, broccoli, brussel sprouts, peppers, apple, pear, peach, apricot, carnations and roses. More specifically, in alternative embodiments, plants for which the invention may be used in modifying fatty acid content include oil crops of the *Cruciferae* family: canola, rapeseed (*Brassica* spp.), crambe (*Crambe* spp.), honesty (*Lunaria* spp.) lesquerella (*Lesquerella* spp.), and others; the *Compositae* family: sunflower (*Helianthus* spp.), safflower (*Carthamus* spp.), niger (*Guizotia* spp.) and others; the *Palmae* family: palm (*Elaeis* spp.), coconut (*Cocos* spp.) and others; the *Leguminosae* family: peanut (*Arachis* spp.), soybean (*Glycine* spp.) and others; and plants of other families such as maize (*Zea* spp.), cotton (*Gossypium* sp.), jojoba (*Simonsia* sp.), flax (*Linum* sp.), sesame (*Sesamum* spp.), castor bean

(*Ricinus* spp.), olive (*Olea* spp.), poppy (*Papaver* spp.), spurge (*Euphorbia*, spp.), meadowfoam (*Limnanthes* spp.), mustard (*Sinapis* spp.) and cuphea (*Cuphea* spp.).

The promoter of the invention may direct expression of the elongase gene in a specific tissue or may be otherwise under more precise environmental or developmental control. Such promoters are referred to here as "inducible" promoters. Examples of environmental conditions that may effect transcription by inducible promoters include anaerobic conditions, elevated temperature, or the presence of light. Examples of promoters under developmental control include promoters that initiate transcription only in certain tissues, such as fruit, seeds, or flowers.

In other embodiments, a nucleic acid disclosed herein may be used for cosuppression or antisense inhibition. Antisense inhibition refers to the production of antisense RNA transcripts capable of preventing or reducing the expression of the target protein. The term "antisense" as used herein refers to a nucleotide sequence in which the sequence of residues is in reverse 5' to 3' orientation in relation to the sequence of residues in a sense strand of a nucleic acid coding sequence. A "sense" or "coding" strand or sequence refers to a sequence that may be transcribed by a cell into an mRNA encoding a protein. An "antisense" sequence will therefore generally have the same sequence as the non-coding strand in a DNA duplex, and an antisense RNA will be complementary to an "antisense" sequence. Co-suppression refers to the phenomenon in which expression of a foreign gene which has substantial homology to an endogenous gene results in the suppression of expression of both the foreign and the endogenous gene.

As used herein, the terms "complementary" or "complementarity" when used in reference to polynucleotides refer to polynucleotides which are related by the base-pairing rules, including the rules governing complementarity between DNA and RNA. For example, the sequence 5'-AGT-3' is complementary to the sequence 5'-ACT-3'.

In another embodiment of the invention, the *KCS2* promoter disclosed herein may be operably linked to agronomically important genes, other than the elongase gene disclosed in the present invention. For example, the *KCS2* promoter may be operably linked to the Bt gene, encoding a protein insecticide, to confer insect resistance on the transformed plant (US Patent No. 5,723,756). The *KCS2* promoter may be operably linked to genes encoding proteins,

which have fungal, bacterial and viral inhibiting effects, such as lysozyme, chitinase, attacin, and cecropin (US Patent No. 5,597,946) to confer fungal/bacterial resistance on transformed plants.

5           A cell, tissue, organ, or organism into which has been introduced a foreign nucleic acid, is considered "transformed", "transfected", or "transgenic". A transgenic or transformed cell or organism also includes progeny of the cell or organism and progeny produced from a breeding program employing a transgenic plant as a parent in a cross and exhibiting an altered phenotype resulting from the presence of a recombinant nucleic acid construct. A transgenic  
10       plant is therefore a plant that has been transformed with a heterologous nucleic acid, or the progeny of such a plant that includes the transgene. The invention provides vectors, such as vectors for transforming plants or plant cells. The term "vector" in reference to nucleic acid molecule generally refers to a molecule that may be used to transfer a nucleic acid segment(s) from one cell to another.

15           Another aspect of the invention provides transgenic plant cells, plant tissues derived from such plant cells, and descendants thereof. Nucleic acids of the invention may be introduced into the genome of the desired plant host by a variety of conventional techniques which include, without limitation, electroporation and microinjection of plant cell protoplasts  
20       and polyethylene glycol precipitation (such as are disclosed in Paszkowski *et al.* (1984); Fromm *et al.*, (1985); Rogers *et al.*, (1986); and in U.S. Patent Nos. 4,684,611; 4,801,540; 4,743,548 and 5,231,019), ballistic methods such as DNA particle bombardment (for example as disclosed in Klein, *et al.*, (1987); Gordon-Kamm, *et al.* (1990); and in U.S. Patent Nos. 4,945,050; 5,015,580; 5,149,655 and 5,466,587); *Agrobacterium*-mediated transformation  
25       methods (such as those disclosed in Horsch *et al. Science* (1984); Fraley *et al.*, (1983); and U.S. Patent Nos. 4,940,838 and 5,464,763). Alternative transformation protocols are disclosed for example in U.S. Pat. No. 5,584,807; 5,501,967; Fraley *et al.* (1982) *Proc. Natl. Acad. Sci. USA* 79:1859-1863; Krens *et al.* (1982) *Nature* 296:72-74).

30           Transformed plant cells, which may be derived by any of the above transformation techniques, may be cultured to regenerate whole plants having the transformed genotype and displaying a desired phenotype, as for altered VLCFA levels. A variety of plant culture techniques may be used to regenerate whole plants, such as described in Gamborg and Phillips (1995); Evans *et al.* (1983); or Binding, (1985); or in Klee (1987).

One of skill will recognize that after the nucleic acid is stably incorporated in transgenic plants and confirmed to be operable, it can be introduced into other plants by sexual crossing. Any of a number of standard breeding techniques may be used, depending upon the species to be crossed.

Nucleic acids of the invention may also be used as a plant breeding tool, as molecular markers to aid in plant breeding programs. Such techniques would include using the gene itself as a molecular probe or using the DNA sequence to design PCR primers to use PCR based screening techniques in plant breeding programs.

The invention now being generally described, it will be more readily understood by references to the following examples, which are included for purposes of illustration only and are not intended to limit the invention unless so stated.

#### Example 1

##### Cloning KCS2

Following discovery and sequencing of the region of KCS2 in *Arabidopsis*, synthetic oligonucleotides homologous to portions of the KCS2 genomic sequence were prepared and used as primers to amplify either 1508 bp or 1001 bp of continuous KCS2 DNA sequence by PCR. As shown in Figure 1, the upstream primer was 5'-GTATCATCAACAAAAATATC-3' (kcs1) in combination with the downstream primer 5'-CAAAGATCGATCTTAACC-3' (kcs2) for the PCR-synthesis of the 1508 bp genomic DNA fragment, which includes a complete coding sequence. Primers 5'-CGATCACGGAGTAGAGAA-3' (kcs3) and 5'-GGACAGTTTCTAAAGCAG-3' (kcs4) were used for the PCR-amplification of the 1001 bp of the 5' untranslated region (promoter fragment). The amplified 1508 bp fragment was subcloned in the EcoRI site of the plasmid pCR2.1 (Invitrogen) to produce plasmid pCR-KCS2, whereas the 1001 bp promoter fragment was subcloned in the SmaI site of the plasmid pGEM7zf (Promega) to produce plasmid pGEM-proKCS2. The cloned DNA fragments were then completely sequenced on both strands using an automatic sequencer by the dideoxy chain termination method. These two fragments together comprise 2508 bp of continuous genomic DNA sequence shown in Figure 1. This DNA sequence represents a 1001 bp 5' untranslated region (i.e., nucleotides preceding the first ATG codon), and a 1461 bp open reading frame that encodes a 487 amino acid protein with a predicted molecular weight of 54,900 Da (KCS2).

## Example 2

### Expression of KCS2 in Yeast

The full length *KCS2* cDNA was expressed in *Saccharomyces cerevisiae* linked to the GAL10 inducible promoter. The *KCS2* gene was cleaved from pCR2.1 with EcoRI and inserted into the EcoRI site of the polylinker of the *S. cerevisiae* expression vector pESC-TRP (Stratagene), resulting in the plasmid pESC-KCS2. The construct was transformed into the *S. cerevisiae* strain YPHY99 by standard procedures (Schiestl and Gietz, 1989) and transformants selected on minimal medium agar plates lacking tryptophan (Ausubel *et al.*, 1995).

When yeast cells harboring the pESC-KCS2 construct were grown in the presence of galactose, 18:0, 20:0, 22:0 and 24:0 fatty acids accumulated (which are not normally present in the YPHY99 yeast stain, and were not detected in cultures of yeast transformed with the pESC plasmid alone). Previous work on FAE1 indicates that the yeast expression system is a predictive model of elongase condensing enzyme activity in plants (Millar and Kunst, 1997). Accordingly, the yeast expression results disclosed herein shows that the *KCS2* gene encodes an elongase (condensing) enzyme capable of mediating biosynthesis of saturated VLCFAs, including 18:0, 20:0, 22:0 and 24:0 fatty acids, in a wide variety of organisms.

## Example 3

### Activity of the Tissue-Specific *KCS2* Promoter

In order to confirm and more precisely delineate *KCS2* expression patterns, 5' flanking sequences from the *KCS2* genomic clone were translationally fused (operatively linked) to the *uidA* reporter gene encoding beta-glucuronidase (GUS). The inserts were cleaved out of pGEM7zf with BamHI and XbaI and directionally subcloned into the corresponding sites of the binary Ti plasmid pBI101 (Clontech), which contains a promoterless GUS gene (Jefferson *et al.* 1987). The pKCS2-GUS fusion construct in pBI101 was introduced into *Agrobacterium tumefaciens* strain GV3101 (Koncz and Schell, 1986) by heat-shock and selected for resistance to kanamycin (50 (g/ml).

The pKCS2-GUS fusion was introduced into *Arabidopsis* using the floral dip method (Clough and Bent, 1998). Screening for transformed seed was done on 50 micrograms/mL kanamycin as described previously (Katavic *et al.*, 1994). The *KCS2* promoter expression pattern was determined using GUS histochemical assays using buffer containing X-GLUC on

whole seedlings, leaves at different stages of development, stems, flowers, siliques and roots on a large number of *pKCS2*-GUS transgenic lines. Wild type plants treated with GUS buffer under the same conditions were used as controls.

5           The expression pattern observed for the *KCS2* promoter in the *pKCS2*-GUS fusion was consistent in all transgenic lines tested. GUS activity was only observed in the anthers and very young emerging leaves (1-2 mm in size). No GUS expression was detected in any of the other tissues or the wild type controls. Expression of *KCS2* in the anthers was confirmed by RNA blot analysis, using the full *KCS2* ORF (Open Reading Frame) as a probe, confirming that the  
10 *Arabidopsis KCS2* promoter mediates the expression of heterologous proteins in a tissue specific manner.

#### Example 4

##### Analysis of the Specificity of the *KCS2* Gene Product

15           Expression of the *KCS2* gene under the control of GAL10, a galactose-inducible promoter in yeast cells expressing *pESC-KCS2* results in the appearance of four extra peaks in GC chromatograms. These peaks correspond to the saturated VLCFAs 20:0, 22:0, 24:0 and 26:0, suggesting *KCS2* condensing enzyme can elongate fatty acids from C18 to C24 in length.

20           To test the ability of *KCS2* to elongate an acyl chain *in planta*, the full *KCS2* coding sequence was placed behind the *FAE1* seed specific promoter. The recipient plant used for the experiment was the mutant CB25, which contains a lesion in the *FAE1* gene, resulting in a truncated *FAE1* protein. Thus, CB25 plants do not make VLCFAs in the seeds, and all the VLCFAs observed would be the product of the *KCS2* condensing enzyme.

25           Gas chromatography analysis was performed on seeds of 50 transgenic lines and compared to CB25 seeds. A number of lines showed a fatty acid profile different from the one observed for CB25 seeds. The most dramatic difference was observed in the levels of mono-unsaturated C20:1 fatty acid, which in some lines shows an increase of almost 100-fold when  
30 compared to the control, followed by C22:1 fatty acid and C20:0 fatty acid, as shown in Table 1.

Table 1.  
Fatty Acid Profile of KCS2 Expression

FA	CB25	5-6	5-15	5-8	6-8	5-10	5-16	6-7
	% area	% area	% area	% area	% area	% area	% area	% area
16:0	12.18	8.93	9.58	8.89	11.70	8.65	9.22	9.44
16:1	0.67	0.41	0.53	0.46	0.77	0.39	0.43	0.61
18:0	2.92	3.69	4.85	3.93	2.92	3.01	3.49	3.49
18:1	27.59	20.86	19.95	14.07	15.16	20.09	25.37	12.52
18:2	33.47	29.90	28.98	27.62	28.83	30.72	30.97	29.06
18:3	22.02	23.12	19.15	21.60	22.12	22.18	23.32	20.67
20:0	0.63	1.13	1.84	2.10	1.35	1.11	0.87	1.82
20:1	0.24	11.16	14.02	19.49	15.73	12.89	5.74	20.45
22:0	0.16	0.19	0.25	0.29	0.23	0.19	0.18	0.25
22:1	0.00	0.45	0.66	1.32	0.10	0.63	0.26	1.55
24:1	0.10	0.14	0.17	0.21	0.17	0.14	0.14	0.14

5

The identity of the peaks was estimated by comparing their retention times to known fatty acid standards.

#### 10 Example 5

##### Utility of KCS2 for Seed Oil Modification

Results of the experiments described above show that the *KCS2* gene is primarily expressed in flower buds and demonstrate that *KCS2* condensing enzyme can elongate VLCFAs in *Arabidopsis* seeds. In an attempt to co-suppress *KCS2* in order to obtain a loss-of-  
 15 function phenotype, as well as a gain-of-function, over-expression phenotype, the strong CaMV 35S promoter was used to constitutively express *KCS2* in all plant tissues.

RNA blot analyses carried out on randomly selected transgenic lines to assess the level of expression of the 35S-*KCS2* transgene demonstrated high levels of expression of the *KCS2*  
 20 gene in a number of lines. To determine whether high levels of the *KCS2* expression translated into higher levels of VLCFAs in the seed, seed of the 35S-*KCS2* plants showing the highest levels of the *KCS2* transcript were subjected to gas chromatography analyses.

Fatty acid profiles of all the transgenic lines tested differ significantly from the wild  
 25 type as shown in Figure 2. The main difference was observed in the level of 20:1 fatty acid (approximately 18% versus 12% in wild type), and a reduced 18:1 content. Apparently, the expression of *KCS2* in seeds increases the rate of conversion of 18:1 to 20:1 fatty acids and as

a result, all the transgenic 35S-*KCS2* lines accumulate significantly more total VLCFAs than the wild type seeds. A calculation of the proportion of VLCFAs with respect to total fatty acids showed that all the transgenic lines tested have over 20% of VLCFAs, whereas in the wild type VLCFAs account for 16.9% of total fatty acids.



**REFERENCES**

All publications, including patent applications referred to herein and including the following, are hereby incorporated by reference:

5 WO 95/15387

WO 96/13582

U.S. Patent No. 4,940,838

10

U.S. Patent No. 5,464,763

U.S. Patent No. 4,945,050

15 U.S. Patent No. 5,015,580

U.S. Patent No. 5,149,655

U.S. Patent No. 5,466,587

20

U.S Patent No. 4,684,611

U.S Patent No. 4,801,540

25 U.S Patent No. 4,743,548

U.S Patent No. 5,231,019

US Patent No. 5,597,946

30

US Patent No. 5,723,756

Ausubel, F.M. *et al.* eds., (1995) Current Protocols in Molecular Biology (New York: John Wiley and Sons).

Altschul S.F. *et al.* (1990) Basic local alignment search tool. *J Mol Biol.* 215(3):403-410.

Binding, "Regeneration of Plants, Plant Protoplasts", CRC Press, Boca Raton, 1985.

- 5 Clough, S.J. and Bent, A.F. (1998) Floral dip: a simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J.* 16: 735-743.

Evans *et al.* "Protoplasts Isolation and Culture", Handbook of Plant Cell Culture, Macmillan Publishing Company, New York, 1983.

10

Fraley *et al.*, *Proc. Natl. Acad. Sci. USA* 80:4803 (1983).

Fromm *et al.*, *Proc. Natl. Acad. Sci. USA* 82:5824 (1985).

Gamborg and Phillips, "Plant Cell, Tissue and Organ Culture, Fundamental Methods",

- 15 Springer Berlin, 1995.

Gordon-Kamm, *et al.* "The Plant Cell" 2:603 (1990).

Haque, M.M. *et al.* (1992) Inheritance of leaf epicuticular wax content in rice. *Crop Sci.* 32:

- 20 865-868.

Harlow and Lane (1988) *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, New York.

- 25 Horsch *et al.* *Science* 233:496 (1984).

Innis, *et al.* eds. PCR Protocols. A Guide to Methods and Application. (Academic Press, Inc. San Diego, CA. 1990).

- 30 James, D.W. and Dooner, H.K. (1990) Isolation of EMS induced mutations in *Arabidopsis* altered in seed fatty acid composition. *Theoretical and Applied Genetics* 80: 241-245.

Jefferson, R.A. *et al.* (1987) GUS fusions: -glucuronidase as a sensitive and versatile gene fusion marker system in higher plants. *EMBO J.* 6: 3901-3907.

Katavic, V. *et al.* (1994) In plant transformation of *Arabidopsis thaliana*. *Mol.Gen.Genet.* 245: 363-370.

Klee *et al.*, *Ann. Rev. of Plant Phys.* 38:467 (1987).

Klein, *et al.*, *Nature* 327:70 (1987).

Koncz, C. and Schell, J. (1986) The promoter of TL-DNA gene 5 controls the tissue-specific expression of chimeric genes carried by a novel type of *Agrobacterium* binary vector. *Mol. Gen. Genet.* 204: 383-396.

Kunst, L. *et al.* (1992) Fatty Acid Elongation in developing seeds of *Arabidopsis thaliana*. *Plant Physiol. Biochem.* 30: 425-434.

Lassner *et al.* (1996) A Jojoba  $\beta$ -keto-acyl-coA-synthase cDNA compliments the Canola fatty acid elongation mutation in transgenic plants. *Plant Cell* 8: 281-292.

Lemieux *et al.* (1990) Mutants of *Arabidopsis* with alterations in seed lipid fatty acids. *Theoretical and Applied Genetics* 80: 234-240.

Millar, A.A., and Kunst, L. (1997) Very-long-chain fatty acid biosynthesis is controlled through the expression and specificity of the condensing enzyme. *Plant J.* 12: 121-131.

Millar, A.A. *et al.* (1999) CUT1, an *Arabidopsis* gene required for cuticular wax biosynthesis and pollen fertility, encodes a very-long-chain fatty acid condensing enzyme. *Plant Cell* 11: 825-838.

Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443.

O'Toole, J.C. and Cruz, R.T. (1983) Genotypic variation in epicuticular wax of rice. *Crop Sci.* 23: 392-394.

Paszkowski *et al.* *EMBO J.* 3:2717 (1984).

Pearson and Lipman, 1988, *Proc. Natl. Acad. Sci. USA* 85:2444.

Post-Beittenmiller, D. (1996) Biochemistry and molecular biology of wax production in plants. *Annu. Rev. Plant Physiol. Mol. Biol.* 47: 405-430.

Rogers *et al.*, *Methods Enzymol.* 118:627 (1986).

Sambrook, *et al.* *Molecular Cloning: A Laboratory Manual.* (2<sup>nd</sup> ed., Vols. 1-3, Cold Spring Harbor Laboratory (1989).

Schiestl, R.H., and Gietz, R.D. (1989) High efficiency transformation of intact yeast cells using single stranded nucleic acids as a carrier. *Curr. Genet.* 16: 339-346.

Smith and Waterman (1981) *Adv. Appl. Math* 2:482.

Tijssen, (1993) *Laboratory Techniques in Biochemistry and Molecular Biology – Hybridization with Nucleic Acid Probes, Part I, Chapter 2 “Overview of Principles in Hybridization and the Strategy of Nucleic Acid Probe Assays”* (Elsevier, New York).

## CONCLUSION

Although various embodiments of the invention are disclosed herein, many adaptations and modifications may be made within the scope of the invention in accordance with the common general knowledge of those skilled in this art. Such modifications include the substitution of known equivalents for any aspect of the invention in order to achieve the same result in substantially the same way. Numeric ranges are inclusive of the numbers defining the range. In the specification, the word "comprising" is used as an open-ended term, substantially equivalent to the phrase "including, but not limited to", and the word "comprises" has a corresponding meaning. Citation of references herein shall not be construed as an admission that such references are prior art to the present invention. All publications, including but not limited to patents and patent applications, cited in this specification are incorporated herein by

reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein and as though fully set forth herein.

What is claimed is:

1. A recombinant nucleic acid molecule comprising a heterologous nucleic acid coding sequence encoding a plant long chain fatty acid condensing enzyme, wherein:
  - 5 a) the nucleic acid coding sequence is derived from an *Arabidopsis KCS2* coding sequence; or
  - b) the plant very long chain fatty acid condensing enzyme catalyses the condensation of malonyl-CoA with a C16, C18, C20 or C22 acyl-CoA, wherein the plant very long chain fatty acid condensing enzyme has an amino acid sequence that is at least 70% identical to an
  - 10 *Arabidopsis KCS2* amino acid sequence when optimally aligned; or
  - c) the nucleic acid coding sequence hybridizes under stringent conditions to a complement of the *Arabidopsis KCS2* coding sequence; or
  - d) the nucleic acid coding sequence at least 70% identical to the *Arabidopsis KCS2* coding sequence when optimally aligned.
- 15 2. The recombinant nucleic acid molecule of claim 1 wherein the nucleic acid coding sequence is derived from the *Arabidopsis KCS2* coding sequence.
3. The recombinant nucleic acid molecule of claim 1 wherein the plant very long chain fatty acid condensing enzyme catalyses the condensation of malonyl-CoA with a C16, C18, C20 or C22 acyl-CoA, wherein the plant very long chain fatty acid condensing enzyme has an amino acid sequence that is at least 70% identical to the *Arabidopsis KCS2* amino acid sequence when optimally aligned.
- 20 4. The recombinant nucleic acid molecule of claim 1 wherein the nucleic acid coding sequence hybridizes under stringent conditions to the complement of the *Arabidopsis KCS2* coding sequence.
- 25 5. The recombinant nucleic acid molecule of claim 1 wherein the nucleic acid coding sequence at least 70% identical to the *Arabidopsis KCS2* coding sequence when optimally aligned.
- 30

6. The recombinant nucleic acid molecule of claim 1 wherein the nucleic acid coding sequence at least 90% identical to a wild-type *Arabidopsis KCS2* coding sequence when optimally aligned.
- 5 7. The recombinant nucleic acid molecule of claim 1 wherein the nucleic acid coding sequence at least 95% identical to a wild-type *Arabidopsis KCS2* coding sequence when optimally aligned.
8. An isolated nucleic acid molecule comprising a nucleic acid coding sequence that  
10 encodes a plant long chain fatty acid condensing enzyme, wherein:  
a) the nucleic acid coding sequence is derived from an *Arabidopsis KCS2* coding sequence; or  
b) the plant long chain fatty acid condensing enzyme catalyses the condensation of  
malonyl-CoA with a C16, C18, C20 or C22 acyl-CoA, wherein the plant very long chain fatty  
15 acid condensing enzyme has an amino acid sequence that is at least 70% identical to an  
*Arabidopsis KCS2* amino acid sequence when optimally aligned; or  
c) the nucleic acid coding sequence hybridizes under stringent conditions to a  
complement of the *Arabidopsis KCS2* coding sequence; or  
d) the nucleic acid coding sequence is at least 70% identical to the *Arabidopsis KCS2*  
20 coding sequence when optimally aligned.
9. The isolated nucleic acid molecule of claim 8, wherein the nucleic acid coding sequence is derived from the *Arabidopsis KCS2* coding sequence.
- 25 10. The isolated nucleic acid molecule of claim 8, wherein the plant long chain fatty acid condensing enzyme catalyses the condensation of malonyl-CoA with a C16, C18, C20 or C22 acyl-CoA, wherein the plant very long chain fatty acid condensing enzyme has an amino acid sequence that is at least 70% identical to an *Arabidopsis KCS2* amino acid sequence when optimally aligned.
- 30 11. The isolated nucleic acid molecule of claim 8, wherein the nucleic acid coding sequence hybridizes under stringent conditions to a complement of the *Arabidopsis KCS2* coding sequence.

12. The isolated nucleic acid molecule of claim 8, wherein the nucleic acid coding sequence is at least 70% identical to the *Arabidopsis KCS2* coding sequence when optimally aligned.

5 13. The isolated nucleic acid molecule of claim 8, wherein the nucleic acid coding sequence is at least 90% identical to a wild-type *Arabidopsis KCS2* coding sequence when optimally aligned.

10 14. The isolated nucleic acid molecule of claim 8, wherein the nucleic acid coding sequence is at least 95% identical to a wild-type *Arabidopsis KCS2* coding sequence when optimally aligned.

15 15. A recombinant nucleic acid molecule comprising a promoter sequence operably linked to a nucleic acid sequence, wherein the promoter sequence is capable of mediating gene expression in anthers and in very young leaves in *Arabidopsis* and:

a) is derived from an *Arabidopsis KCS2* promoter sequence; or

b) hybridizes under stringent conditions to the *Arabidopsis KCS2* promoter sequence;

or,

20 c) is at least 70% identical to the *Arabidopsis KCS2* promoter sequence when optimally aligned.

16. The recombinant nucleic acid molecule of claim 15, wherein the promoter sequence is derived from the *Arabidopsis KCS2* promoter sequence.

25 17. The recombinant nucleic acid molecule of claim 15, wherein the promoter sequence hybridizes under stringent conditions to the *Arabidopsis KCS2* promoter sequence.

18. The recombinant nucleic acid molecule of claim 15, wherein the promoter sequence is at least 70% identical to the *Arabidopsis KCS2* promoter sequence when optimally aligned.

30 19. The recombinant nucleic acid molecule of claim 15, wherein the promoter sequence is at least 90% identical to a wild-type *Arabidopsis KCS2* promoter sequence when optimally aligned.



20. A nucleic acid probe comprising a probe sequence that:
- a) hybridizes under stringent conditions to a portion of an *Arabidopsis KCS2* genomic sequence; or
  - b) is at least 70% identical to the portion of an *Arabidopsis KCS2* genomic sequence when optimally aligned.
21. The nucleic acid probe of claim 20 wherein the probe sequence hybridizes under stringent conditions to a portion of the *Arabidopsis KCS2* genomic sequence.
22. The nucleic acid probe of claim 20 wherein the probe sequence is at least 70% identical to the portion of the *Arabidopsis KCS2* genomic sequence when optimally aligned.
23. The nucleic acid probe of claim 20 wherein the probe sequence is at least 90% identical to a portion of a wild-type *Arabidopsis KCS2* genomic sequence when optimally aligned.
24. A transgenic plant comprising the recombinant nucleic acid molecule of any one of claims 1 through 7.
25. A part of the transgenic plant of claim 24.
26. The part of the transgenic plant of claim 25, wherein the part is a seed.
27. The transgenic plant of claim 24, wherein the transgenic plant has a modified phenotype compared to a non-transgenic plant of the same species.
28. A transgenic cell comprising the recombinant nucleic acid molecule of any one of claims 1 through 7.
29. The transgenic cell of claim 28, wherein the cell is a plant cell.
30. A method of producing a transgenic plant comprising introducing into the plant the isolated nucleic acid molecule of any one of claims 8 through 14.

31. A progeny plant produced by sexual or asexual propagation of the transgenic plant produced by the method of claim 30.

32. A purified protein encoded by the recombinant nucleic acid molecule of any one of claims 1 through 7.

33. A recombinant vector comprising the recombinant nucleic acid molecule of any one of claims 1 through 7.

34. A recombinant antisense nucleic acid molecule wherein a portion of the heterologous nucleic acid coding sequence of claim 1 is in reverse orientation relative to an adjacent promoter sequence.

35. The recombinant antisense nucleic acid of claim 34, wherein the recombinant antisense nucleic acid encodes an antisense RNA that:

a) hybridizes under stringent conditions to a complement of a portion of the *Arabidopsis KCS2* coding sequence; or

b) is at least 70% identical to a portion of the *Arabidopsis KCS2* coding sequence when optimally aligned.

36. A Transgenic plant or plant cell comprising the recombinant antisense nucleic acid of claim 34 or 35.

37. A method of isolating a nucleic acid molecule encoding a plant long chain fatty acid condensing enzyme, the method comprising hybridizing a nucleic acid preparation with the nucleic acid probe of any one of claims 20 through 23.

Figure 1

```

1   tgttgtggag gacttgtgag aaccaccacc agagtccgac atcgcgatca
51  cggaagtagag aaagtcaaaa ctacttctct cagacggatt agtttggttt
      kcs3->
101 gctggagatt gttccaagaa agagaaacgt taggagcaaa caaacaanaag
151 agaaaagacg atgatgactg atgagagctt taacaaaaaa ataaaatgag
201 agagctcaac gggtagaatt gtgagacttg agagagtgtt tcctatttaa
251 ggcattgcgat tagtgtttat taccgagaatg ccaccgaacg agtacatatt
301 aatgtatagt atgttaatga tagtctaact aaaatttggg ttttattgaa
351 atagaatttt gtaagaataa tgaggatctg taatatagct ggatttgcat
401 taaatcgtag gccgttggtg atcgaaatta gttaataaaa tgttttagca
451 tataatgttg gtgcttccga catgtttatt ggacaataat accatatttt
501 ttctttggga tcttaaaaaa attgaggaag aaaatagtaa aatagtcaaa
551 cttagggttac atcataatgg gccatttctt tgagttgtga ttgatctcca
601 aagatataca tagatttaca caagatcaaa agaaaaacaa ttgggcctaa
651 accccaagcc catatcaacg tccattatca ttaagattcc ttttttctt
701 gaaatttgaa aatttgaaat tcgattcaaa tctactctct ctgttttttt
751 cccataaaaa tctgaaaaac cagaagcttc ttcactcact ttcctcttga
801 tatcttccat tagttggccg atacacatga cgccaaatac atcaatggcg
851 actcttctct gttttttagt tatatcaaac tcccacccaa cctgcagaag
901 aaaaaatggt gtctataaac acatcccctt acgatttctt ctctatctct
951 ctcacagtat ctatatatac gcacacaaac ccagattcag tttctcatca
1001 gtatcatcaa caaaaatctc aaagattctg ctttagaana tgtccATGga
      kcs1 ->          <- kcs4
1051 tgctaattgga ggacctgtac agatccggac ccaaaactac gtcaagcttg
1101 gttatcacta tctgatcact cactttttta aactcatgtt cctccctcta
1151 atggctgttt tgttcatgaa tgtctcattg ttaagcctaa accatcttca
1201 gctctattac aattccaccg gattcatctt cgtcattact ctgccattg
1251 tcggatccat tgtcttcttc atgtctcgac ctagatccat ctaccttcta
1301 gattactctt gctacctccc gccttcgagt caaaaagtta gctaccagaa
1351 attcatgaag aactctagtt tgattcaaga tttcagcgaa acttctcttg
1401 agttccagag gaagatcttg attcgctctg gtctcgggtg agagacttat
1451 ttaccggatt ctattcactc tatccctccg cgtcctacta tggctgcagc
1501 gcgtgaagaa gcggagcagg taatcttcgg tgactcgcac aatcttttcg
1551 agaatacaaa aatcaatcct agggagattg gtgttcttgt tgtgaattgt
1601 agtttgttta accctacgcc ttctttatcc gccatgattg ttaacaagta
1651 taagcttaga ggaaacatta agagctttta ccttgaggga atgggatgta
1701 gtgctgggtg tatcgcggtg gatctagcta gtgatatgtt acaaatccat
1751 aggaacactt ttgctcttgt ggtagtact gagaacatca ctcagaattg
1801 gtatttttgt aacaagaaaag caatgttgat ccctaattgc ttgtttagag
1851 ttggtgggtc cgcggttctg ctttcgaaca agcctttgga tcgaaaacga
1901 tccaagtata agcttggttca tacggtcagg actcataaag gatctgatga
1951 gaacgcattc aattgtgtgt atcaagaaca agatgagtgt ttgaaaaccg
2001 gagtttcttt gtctaaagat cttatggcta tagctggaga agctttaaag
2051 acgaatatca cttctttggg tcctctggtt cttcctataa gcgagcagat
2101 tctgttcttt gcgacttttg ttgctaagag attgttcaat gacaagaaga
2151 agaagcctta cataccggat ttcaagcttg ctttagatca tttctgtatt
2201 cacgcgggag gtagagccgt gattgatgag ctagagaaga gtttaaagct
2251 ttctccaaaa catgttgagg cgtctagaat gactttgcat agatttgaa
2301 acacttcttc tagctctata tggatgaat tggcttacac ggaagctaaa
2351 ggaagaatga ggaaaggaaa cagagtttg cagattgctt ttggtagcgg
2401 gtttaagtgt aacagcgcgg tttgggtggc tcttcgcaat gtcgagccct
2451 cggttaacaa tccttgggaa cattgcattc atagatatcc ggttaagatc
      <- kcs2
2501 gatctttga

```